

ORIGINAL ARTICLE

Socioeconomic status and race as social determinants of health to be considered in clinical use of pre-season vestibular and oculomotor tests for concussion

Jessica Wallace¹, Phillip Worts^{2,3,4}, Ryan Moran¹, Justin Mason⁵, Katherine K. Weise⁶, Mark Swanson⁶, Nicholas Murray⁷

1. University of Alabama – Department of Health Science, 270 Kilgore Lane, Capital Hall 2106, Tuscaloosa, AL 35405

2. Tallahassee Orthopedic Clinic 3334 Capital Medical Blvd Suite 600 Tallahassee, FL 32308

3. Florida State University - Department of Nutrition, Food and Exercise Sciences, Tallahassee, FL 32308

4. Florida State University Institute of Sports Sciences & Medicine Tallahassee, FL 32308

5. University of Florida – Department of Occupational Therapy, 1225 Center Drive, Room 2119 Gainesville, FL 32611

6. University of Alabama-Birmingham School of Optometry, 1716 University Blvd, Henry Peters Building 506, Birmingham, AL 35233

7. University of Nevada, Reno – School of Community Health Sciences, 1664 N. Virginia Street m/s 0274, Reno, NV 89557

*Corresponding author

Jessica Wallace

University of Alabama – Department of Health Science 270 Kilgore Lane, Capital Hall 2106 Tuscaloosa, AL 35405

Tel: (o) 205-348-4577 / (f) 205-348-7568

Email: Jswallace1@ua.edu

Article information:

Received: June 2, 2020

Revised: August 8, 2020

Accepted: September 24, 2020

Abstract

Background: Aside from racial and socioeconomic disparities in computerized neurocognitive testing and symptomology, there is a scarcity of research representing more diverse populations on other widely used tests for concussion, including vestibular and visual assessment.

Aim: To investigate if racial and socioeconomic differences exist on baseline Vestibular Ocular Motor Screening (VOMS) and King-Devick (K-D) test performance in high school student-athletes.

Methods: A total of 670 participants (66.1% White, 33.9% Black) with a mean age of 15.43 ± 1.2 years were administered a baseline VOMS, average NPC distance, and K-D test. The exposure variables included race (White or Black) and socioeconomic status (SES), defined as free-and-reduced lunch status (FRL or No-FRL). FRL status was determined by each participant's school SES. The outcome variables consisted of baseline VOMS item symptom provocation scores, average NPC distance and K-D baseline time. A series of Mann-Whitney *U* tests were performed for K-D baseline time, NPC distance, and VOMS items with FRL status or race as a between subject factor. Two multivariable linear regressions were run to assess the association of (1) K-D baseline times using FRL, race, sex, and corrected vision as variables in the model and (2) average NPC distance using FRL, race, sex, and corrected vision as variables in the model.

Results: When adjusting for multiple comparisons, FRL athletes had slower (worse) K-D times ($p < .001$) than non-FRL athletes. Black athletes had significantly lower mean NPC distance compared to White athletes at baseline ($p = .02$) and FRL status athletes reported a significantly greater (worse) mean symptom provocation following the visual motion sensitivity (VMS) item on the VOMS ($p = .02$); however, these findings were no longer significant following adjustments for multiple comparisons. No differences were noted for any remaining VOMS items. The first model explained 3.9% of the total variance of K-D baseline times whereas the second model was not significant.

Conclusions: Racial and SES differences existed on average NPC distance, and the K-D test at baseline. Possible explanations for group differences may be neurobiological, anatomical, and/or disparity in nature. With a higher probability of undiagnosed and uncorrected vision impairment,

vestibular dysfunction, and saccadic eye tracking deficits likely to be more apparent as a consequence of poverty or health inequities, it is important that healthcare providers, especially those that diagnose and treat concussions, understand that performance on the VOMS and K-D tests at baseline may be subject to sociodemographic factors of SES and race.

Relevance for Patients: To provide the most culturally competent care, clinicians should consider sociodemographic variables of race and SES as social determinants of health worthy of attention on objective and subjective measures of baseline concussion assessment.

Key Words: race, socioeconomic status, concussion, VOMS, vestibular, King-Devick

1. Introduction

A concussion occurs following a direct or indirect traumatic force that is transmitted to the brain, which initiates pathophysiological processes (i.e., neurometabolic cascade) and results in transient brain dysfunction [1,2]. Patients may experience immediate or delayed onset of symptoms including headache, dizziness, and fogginess, abnormal vestibular function, altered behavior, impaired cognition, sleep dysregulation, cardiac autonomic dysfunction, or visual dysfunction [3–5]. The diagnosis of a concussion is centered around a multifaceted assessment battery which includes symptoms, neurocognition, postural stability and vision/vestibular assessments [1]. Concussion recovery may vary by age and prior history of concussion [6]; however, most adolescent concussions are believed to resolve in ≤ 4 weeks [3]. Unfortunately, a multitude of sociodemographic factors may affect clinical concussion diagnosis, treatment, and outcomes, which may be overlooked by clinicians.

Socioeconomic status (SES) (also known as socioeconomic position, sociodemographic characteristics, or sociocultural factors) has historically influenced clinical outcomes, dating as far back as the Pellegra epidemic in the 1900s [7,8]. Specific to concussion management, researchers are still reporting disparities in healthcare a century later, which disproportionately affects low-income patients and certain racial groups. Cognitive test performance [9], access to athletic trainers in secondary schools [10,11], access to specialized concussion care [12], and prolonged hospitalization

following traumatic brain injuries are areas of inequity that warrant attention and must be addressed [13]. School SES, measured as a Title I classification [14], is a poverty indicator and the receiving of federal subsidies; however, these funds may not be prioritized for student-athlete health and safety. Further, 54% percent of high schools in the United States (US) meet school-wide Title I status and 52% of high school students in the US are eligible for free and reduced-priced lunch (FRL) [15]. This proportion highlights the number of children nationwide who likely represent a lower SES. Many youth of low SES or a racial/ethnic minority are medically uninsured or insured by Medicaid [16]. This may limit their ability to seek specialized care for concussion. In fact, racial and ethnic minority youth sport participants are less likely to be diagnosed with a concussion in the emergency department compared to White youth [17]. Considered collectively, socioeconomically disadvantaged patients and those of non-white racial groups may not be positioned to attain optimal clinical outcomes if they suffer a concussion. Additionally, if socioeconomically disadvantaged patients do seek medical attention following a concussion, it is uncertain if clinical diagnostics are capable of quantifying the influence of SES status on test results.

During various cognitive tests, processing speed [9,18–21], memory [9,18,20–22], and reaction time [18,20] have all shown to be susceptible to performance differences amongst racial groups at baseline or post-injury. Additionally, total symptom reporting was significantly higher in Black athletes versus White athletes [23] and Black children reported higher cognitive-related symptoms post-concussion than their White counterparts [24]. Similarly, low SES groups performed worse on processing speed, memory, and reaction time tests and reported higher symptom scores [25]. Aside from racial and socioeconomic disparities in computerized neurocognitive testing and symptomology, there is a paucity of research representing more diverse populations on other widely used tests for concussion, including those that are designed to assess vestibular and visual dysfunction.

Concussion tests gaining rapid traction and clinical utilization, such as the King-Devick (K-D) Test [26] or Vestibular/Ocular Motor Screening (VOMS) [27], have been used to assist clinicians in the detection of vestibular-ocular impairment following a suspected concussion, yet little is known about racial and SES differences on test performance of the K-D and VOMS. However, on standard

vision tests, there have been differences reported in prevalence rates of abnormal findings across racial groups [28–30]. A higher percentage of Black (14.1%) and Hispanic (14.2%) children failed one or more vision screening tests when compared to White children (11.0%) [28]. Likewise, the National Health and Nutrition Examination Survey (NHANES) estimates suggested that visual impairment was 2-3 times less prevalent in participants who reported a poverty income ratio that was more than two times above the poverty level versus their lower-income counterparts [31]. Further, increased odds of vestibular dysfunction were reported when comparing Black and Mexican American races/ethnicities to Whites [32]. Considering the VOMS and K-D both include vision performance metrics [e.g., Near Point of Convergence (NPC) in the VOMS and saccadic tracking in the K-D], the aforementioned clinical data would suggest that race and SES are worthy of considering when interpreting test data during a concussion evaluation. Therefore, the purpose of the current study was to investigate if racial and/or socioeconomic differences exist in performance on VOMS items, including average NPC distance, and K-D test at baseline in high school student-athletes.

2. Methods

2.1 Design:

This study utilized a cross-sectional design. The exposure variables were race (White or Black) and socioeconomic status, defined as free-and-reduced lunch status (FRL or No-FRL). FRL status was determined by each participant's school SES: Title I or Non-Title I. Title I school SES was defined as high poverty which includes schools where a greater percentage of students in attendance qualify for FRL, and non-Title I school SES was defined as low poverty included schools of which a greater percentage of students in attendance did not qualify for FRL [14]. Title I status was confirmed with administrators at each school district. The outcome variables consisted of baseline VOMS item symptom provocation scores, average NPC distance and K-D baseline time.

2.2 Participants:

A total of 868 high school student-athlete participants from 10 high schools in Northeast Ohio initially enrolled in the study. A total of 670 participants between the ages of 13 and 18 years (15.43 ± 1.2 years) met the study's inclusionary criteria. Any student-athlete diagnosed with a self-reported learning disability, concussion within the past 3 months or any athlete who was missing baseline data on one of the two outcome measures ($n=99$) was excluded from the study. Additionally, individual's whose self-reported race was not White or Black (e.g., Hispanic/Latino, Asian, American Indian, Pacific Islander) were excluded from the analysis ($n=99$) due a low comparative sample size of additional minority races and ethnicities. A breakdown of participant demographics is provided in Table 1.

Table 1: Participant Demographics

	FRL Status		
	No FRL (N=232)	FRL (N=438)	Total (N=670)
Race			
White	202 (87.1%)	241 (55.0%)	443 (66.1%)
Black	30 (12.9%)	197 (45.0%)	227 (33.9%)
Sex			
Female	59 (25.4%)	87 (19.9%)	146 (21.8%)
Male	173 (74.6%)	351 (80.1%)	524 (78.2%)
Prescribed Glasses or Contacts			
Yes	113 (48.7%)	187 (42.7%)	300 (44.8%)
No	119 (51.3%)	250 (57.3%)	369 (55.2%)
Tested with Corrective Lens			
Yes	69 (29.7%)	98 (22.4%)	167 (25.0%)
No - Not prescribed	119 (51.3%)	251 (57.3%)	370 (55.2%)
No - Did not adhere to prescription	44 (19.0%)	89 (20.3%)	133 (19.9%)
Age			
13	2 (0.9%)	13 (3.0%)	15 (2.2%)
14	48 (20.7%)	129 (29.5%)	177 (26.4%)
15	54 (23.3%)	95 (21.7%)	149 (22.2%)
16	70 (30.2%)	109 (24.9%)	179 (26.7%)

17	52 (22.4%)	83 (18.9%)	135 (20.1%)
18	6 (2.6%)	9 (2.1%)	15 (2.2%)

Data reported as N (%)

2.3 Outcome Measures:

Vestibular Ocular Motor Screening (VOMS) Tool: The VOMS was used to screen each student-athlete on 8 individual items including 1) smooth pursuits, 2) horizontal saccades, 3) vertical saccades 4) convergence 5) near point convergence distance (NPC), 6) horizontal vestibular-ocular reflex (H-VOR), 7) vertical vestibular-ocular reflex (V-VOR), and 8) visual motion sensitivity (VMS). Prior to the assessment, student-athletes reported a baseline rating for headache, dizziness, fogginess, and nausea on a scale of 0 (none)-10 (severe). After each item assessment, student-athletes rated each of the 4 symptoms again. Each VOMS item was administered to see if each subscale provoked any of the previously mentioned four symptoms. Symptom change scores were utilized for analyses and calculated by subtracting the pre-assessment symptom ratings from the post-assessment symptom ratings, to reflect true symptom provocation of each item. NPC was collected using a tape measure and a Bernell fixation device. A total of three trials were collected by having each participant self-report diplopia or exophoria, measuring the distance with a tape measure, and then averaged. Normal NPC values are within 5cm or less for children [33]. The averaged NPC values were utilized for analyses. High internal consistency has been previously reported for baseline NPC distance and K-D times in an adolescent student-athlete sample [34].

King-Devick (K-D) Test: The K-D test is a rapid, number-naming tool requiring micro-saccadic eye movements and reaction time. The K-D test measures the time it takes to read a series of numbers on a series of three test cards as quickly as possible, without making any errors. Scoring is calculated as the cumulative time that it takes to read three test cards, which get increasingly more difficult with each card, error-free. Each participant completed the K-D test twice, and the fastest of the two trials was then recorded as the participant's baseline time. If a participant made an error on one of the 2 trials, he or she completed an additional trial(s) until an error-free attempt was completed. Each test was initiated with uniform directions as provided on the K-D test-card booklet followed by completion of

the 1 demonstration card. High internal consistency (Cronbach $\alpha=0.92$) has been recorded for baseline K-D times in youth populations [35]. Within this study, it is noted that the spiral-bound test card Version 1 of the K-D was used.

2.4 Procedure:

Institutional Review Board approval was granted prior to any data collection. Further, all participants provided parental written consent prior to the start of the study. Participants were recruited from 10 high schools in the Northeast Ohio region by the principal investigator and athletic trainers employed at each high school. During pre-season baseline testing, student-athletes were individually administered the VOMS and K-D tests in a quiet classroom at their respective school. Each student-athlete was tested individually with a trained research team member. Data were collected July 2016 through December 2018.

2.5 Statistical Analyses:

Prior to analysis, the dependent variables were screened for normality (i.e., skewness >2 , kurtosis >9 , Shapiro-Wilk $p < .05$, and Q-Q plot observation). The Shapiro-Wilk test results indicated that the VOMS symptom provocation scores, NPC distance, and K-D test times ($p < .01$) did not follow a normal distribution. Thus, a series of Mann-Whitney U tests were run to assess group differences between race (Black or White) and FRL status (FRL, no-FRL) for pre-test and symptom provocation (i.e., change scores) on individual VOMS items along with NPC distance and K-D baseline times. Prior to modeling, outliers (>3 SDs) for K-D baseline time ($n=9$) and NPC distance ($n=14$) were removed. Two multivariable linear regressions were run to assess the association of (1) K-D baseline times using FRL, race, sex, and corrected vision as variables in the model and (2) average NPC distance using FRL, race, sex, and corrected vision as variables in the model. Due to autocorrelation (assessed via the Durbin Watson statistic), modeling was performed with general least squares and Nagelkerke's pseudo R^2 was reported [36]. Variable selection was informed by research noting that sex differences [37], and the use of corrective lenses, if prescribed, should be worn during the K-D test [38]. Lastly, race may play a factor in prescription and use of corrective lenses [39,40];

thus, it was important to examine the potential association between these factors for both analyses. An alpha level of 0.05 was set *a priori* and *p*-values were adjusted to control for multiple comparisons using the Benjamini-Hochberg procedure [41]. Variables with *p*-values significant after correction were indicated with an asterisk. All analyses were conducted in RStudio (RStudio, Boston, MA) with R version 4.0.2, using the nlme package.

3. Results

3.1 Vestibular/Ocular Motor and K-D Baseline Time Differences

A series of Mann Whitney *U* tests revealed small differences consistent with random sampling between race on pre-test symptoms of the VOMS ($U=49571$, $p=.41$, Cohen's $d=.01$) or symptom provocation scores of any VOMS item (U range= 48867 – 50252, p range = .06 - .97); however, results were not statistically significant (Table 2). Regarding FRL status, no differences were observed between the FRL and non-FRL groups on pre-assessment symptoms of the VOMS ($U=50642$, $p=.11$, Cohen's $d=.14$). Further, no statistically significant differences were observed on the VMS VOMS item ($U=49152.0$, $p=.02$, Cohen's $d=.11$), with the FRL group reporting greater (worse) symptom provocation ($0.10 \pm .9$), than non-FRL ($0.06 \pm .6$). However, this VMS finding was no longer significant following the Benjamini-Hochberg procedure. There were no other statistically significant differences between symptom provocation change scores of any VOMS item between FRL status (U range = 49411 – 50664, p range = .11 - .77). Average NPC distance differed between race ($U=55446$, $p=.02$, Cohen's $d=.16$) but did not differ between the FRL and non-FRL groups ($U=48584.0$, $p=.31$, Cohen's $d=.04$). See Figure 1. Refer to Table 2 for means and standard deviations. All medians and interquartile ranges were 0.00 for VOMS symptoms and NPC distance in the sample and between groups.

The results of the Mann Whitney *U* tests indicated that there was a significant difference between K-D baseline time and FRL ($U = 42513$, $p<.01$, Cohen's $d=.32$), where those who received FRL ($50.19 \pm 8.33s$) had slower (worse) overall K-D times compared to those not receiving FRL (48.19 ± 7.70). Regarding potential racial differences, results of the Mann Whitney *U* tests indicated

that there was not a significant difference between K-D baseline time and race ($U = 51685$, $p = .55$, Cohen's $d = .04$). See Figure 2.

Table 2. Comparison of VOMS Pre-Assessment and Symptom Provocation Change Scores between Race and FRL Status

VOMS Item	White	Black	<i>p</i>	FRL	No FRL	<i>p</i>
Pre-test	0.19 ± 1.3	0.19 ± 0.9	.42	0.25 ± 1.5	0.07 ± 0.5	.11
Smooth Pursuits	0.02 ± 0.3	0.04 ± 0.3	.61	0.03 ± 0.3	0.01 ± 0.1	.73
Horizontal Saccades	0.04 ± 0.4	0.07 ± 0.5	.14	0.06 ± 0.5	0.03 ± 0.2	.60
Vertical Saccades	0.08 ± 0.6	0.11 ± 0.6	.44	0.10 ± 0.7	0.07 ± 0.5	.77
Convergence	0.05 ± 0.5	0.04 ± 0.3	.68	0.04 ± 0.5	0.05 ± 0.4	.71
Horizontal VOR	0.08 ± 0.6	0.08 ± 0.5	.97	0.08 ± 0.6	0.08 ± 0.5	.70
Vertical VOR	0.08 ± 0.6	0.05 ± 0.4	.96	0.05 ± 0.5	0.09 ± 0.5	.31
VMS	0.09 ± 0.6	0.20 ± 1.1	.06	0.10 ± 0.9	0.06 ± 0.6	.03
Average NPC distance (cm)	1.82 ± 3.7	1.27 ± 2.3	.02	1.59 ± 3.2	1.72 ± 3.3	.32
K-D baseline time	49.60 ± 7.9	49.24 ± 8.7	.55	50.19 ± 8.3	48.19 ± 7.7	<.001*

Note: pre-test are non-provocation symptom tasks.

* Significant after Benjamini-Hochberg procedure with false discovery rate 0.05.




Figure 1: Violin plots of free reduced lunch (no or yes) by race and NPC distance (centimeters). Longer times indicate worse performance.




Figure 2: Violin plots of free reduced lunch (no or yes) by race and K-D baseline times (seconds). Longer times indicate worse performance.

3.2 Multivariable Linear Regressions

There was dependence of residuals, as assessed by a Durbin-Watson statistic of 0.368. Thus generalized least squares was used and results indicated that the model to investigate the association of K-D baseline times using FRL, sex, race, and corrected vision as predictors was significant ($F(4,661)=8.94$, Nagelkerke's $R^2=.039$, $p<.001$, adjusted $p<.001$) with FRL ($p<.001$), race ($p=.01$), and sex ($p<.01$) being the significant predictors. There was no evidence of multicollinearity, as assessed by VIF (VIF range 1.00-1.21). The overall model was able to explain 3.9% of the total variance of the dependent variable. See Table 3.

There was dependence of residuals, as assessed by a Durbin-Watson statistic of 0.008. Thus generalized least squares was used and results indicated that the model to investigate the relationship of average NPC distance using FRL, race, sex, and corrected vision was not significant ($F(4,656)=0.73$, Nagelkerke's $R^2=.009$, $p=.20$, adjusted $p=.33$), with race (95% CI= -3.58, -1.99, $p=.03$) being the only significant predictor. There was no evidence of multicollinearity, as assessed by VIF (VIF range 1.00-1.21). See Table 3.

Table 3. Multivariable Linear Regression Results for K-D Baseline Time and Average NPC Distance Measures (# Observations = 669 for both models)

	B	SE	95% CI	p
K-D Baseline Time				
Intercept	46.89	1.91	[43.15, 50.64]	
FRL	3.32	0.77	[1.80, 4.83]	<.001*
Race	-1.99	0.81	[-3.58, -1.99]	.01*
Sex	2.51	0.88	[.79, 4.23]	<.01*
Corrected Vision	-0.31	0.44	[-1.17, 0.56]	.49
Average NPC Distance				
Intercept	2.31	0.72	[0.91, 3.72]	
FRL	0.09	0.29	[-.48, 0.66]	.76
Race	-0.68	0.30	[-1.27, -0.68]	.03*
Sex	0.30	0.33	[-0.34, 0.95]	.35
Corrected Vision	0.15	0.17	[-0.48, 0.17]	.35

*Significant if the 95% CI does not include 0.00 and after Benjamini-Hochberg procedure with false discovery rate <0.05

4. Discussion

The purpose of the current study was to investigate racial and socioeconomic differences in symptom provocation scores on VOMS items, average NPC distance, and K-D test total time at baseline in high school student-athletes. The hypothesis that there would be no differences between groups must be rejected. The primary findings of this study indicate that baseline administration of the VOMS and the K-D test exhibit racial and socioeconomic differences. First, the K-D baseline times were slower (worse) among those that receive FRL ($50.19 \pm 8.3s$) when compared to those that do not receive FRL ($48.19 \pm 7.7s$); however, no significant differences were noted by race. Further, a multivariable model to identify the association between K-D baseline time and independent variables suggests that FRL status and race are both independently associated with K-D baseline time when controlling for other independent variables. As a group, White individuals ($1.82 \pm 3.7cm$) had greater (worse) average NPC distances when compared to Black individuals ($1.27 \pm 2.3cm$) with a small effect size; however, when controlling for multiple comparisons, that difference was no longer significant. Greater symptom provocation reported on the VMS item was noted among those within the FRL group, but again that difference diminished after multiple comparison analyses. It should be noted that average NPC distance was not outside the clinically accepted normal values for the potential diagnosis of concussion [27,43]. Possible explanations for group differences may be neurobiological, anatomical, and/or disparity in nature.

Upwards of 50% of the brain's network is dedicated to vision [44] and adding a vision-based performance measure has enhanced the detection of concussion [45]. Two central findings from this study include a component of vision that appear to be deficient in high school student-athletes. First, a potential saccadic tracking skill deficiency was noted by slower K-D performance among those of FRL. Second, although within a clinically normal range, a potential racial difference in average convergence distance was noted. Both convergence insufficiency and deficient saccadic tracking skills are common binocular vision impairments and the rates of visual impairment are higher in 12-19 years old compared to other age groups, apart from those 60 or older [31].

Although a racial difference in average NPC distance was noted before multiple comparisons, the effect size was small. Black student-athletes had a lower NPC distance and, similarly, black children have been shown to report fewer eye-related symptoms on the Convergence Insufficiency Symptom Survey (CISS) compared to White children [46]. These findings may allude to discrepancies or disparities in objective versus subjective measures for vision among racially diverse student-athletes that warrant further research. Within this study, a difference of approximately 0.6cm is likely clinically insignificant and within normal limits of measurement error. However, these modest racial differences could be explained by high potential for error of objective NPC distance using the nose as the measurement landmark specified with the VOMS. Further, NPC distance using the tip of the nose adds subjectivity to the measure and an influx of anthropometric errors can arise from anatomical variation in nose shape and length. Facial morphology studies using three-dimensional images have demonstrated racial differences in facial structure, including nose length between African Americans and the Welsh. African Americans displayed a less prominent nose and chin and more protrusive nasolabial fold when compared to the Welsh [47]. Currently there is a lack of racial or ethnic diversity in published VOMS norms, yet there is a substantial Black/African American presence in sport, and higher-risk concussion sports. Suggestions to modify facial landmarks used for VOMS NPC measurements include using a less morphologically variable facial landmark, such as the lateral canthi (lateral confluence of upper and lower eyelid margins). Furthermore, assuring standardized administration methods may affect measurements [48]. Adding a more homogenous NPC measurement landmark to the VOMS like the lateral canthus may better align with measures used in eye care practices.

Previous research suggests that rates of uncorrected vision impairment are higher in low income children due to poorer access to eye care services [39,40]. Further, hyperopia is suggested to be higher in non-Hispanic White children as compared to Black children [30]; and, uncorrected hyperopia has been negatively associated with progressive reading skills in youth [49] and adolescents [50] longitudinally. Binocular coordination of eye movements is essential for reading skills that also include the reading of numbers, as required by the K-D. Uncorrected vision problems can severely

impact reading ability, and saccadic tracking skill deficits have been suggested to be a risk factor for poorer reading ability in adolescents [50,51]. Moreover, underdiagnosed and undertreated vision problems that influence reading ability and academic performance often present as frequent eye rubbing or blinking, a short attention span, exo/esophoria, diplopia, losing one's place when reading, tilting the head to one side when reading, or holding reading materials closer to the face [52]. As a vision-based tool that has gained traction in concussion detection, the K-D requires intact saccades and rapid-reading abilities; thus, the poorer performance among those of low SES can, in theory, be explained by higher potential for uncorrected vision impairments.

In addition to possible vision impairments leading to K-D reading deficits, the K-D was also suggested to be a cognitive-related tool due to pre-frontal cortex processing [53]. Rapid-reading ability can also be a consequence of reading skill level, which is linked to academic achievement differences between high SES and low SES groups. Reading skill ability and academic achievement gaps are widened by SES inequities and disparities determined by differences in the kind of school and classroom environment students have access to, resources within the home and neighborhood, and exposure to social capital necessary for success in school [54]. FRL is a widely-used proxy for SES, but SES is a stronger predictor of academic achievement for White students than students of a racial minority [54]. Moreover, the overall slower time to complete the K-D test could be also be due to potential reduced executive functioning and cognitive maturity of those with lower SES [55], as well as a greater likelihood of learning disabilities going undetected and undiagnosed among disadvantaged youth [56]. Results of our study parallel those by Weise et al that found school-type differences on K-D performance among adolescents attending private versus public schools; however, school-type differences in the aforementioned study did not include a measure of SES such as FRL or Title I status [57]. In addition, the literature is sparse regarding how race may influence cognitive maturity or saccadic eye movements; however, the more complex the language [58] and the development of attentional capacity at a young age could influence eye movements and cognition in low-resourced areas [59]. However, these observations are purely speculative given that the K-D test is not a direct measure of saccadic eye movements or cognitive function. It should also be noted that the significant

differences in K-D test baseline results between FRL groups in this study are not believed to be clinically meaningful based on not exceeding the minimal detectable change threshold of 6.10s [60] or the smallest reliable change index threshold of 3.61s [61].

The VOMS symptom provocation score is an entirely subjective measure which is calculated by subtracting the reported score following each item by the pre-test value. This would lead the authors to believe that symptom provocation (i.e.) change scores were therefore subjected to the same limitations of previous symptom reporting tools [62–66]. Two main differences are that many scales use a 0 to 6 Likert scale while the VOMS uses a 0 to 10 Likert scale and the VOMS is designed to look at the change in symptoms to try an account for the individuality of reporting. Specific to race, black student-athletes reported higher symptoms when compared to white student-athletes on a concussion symptom survey [23]; however, Black children have reported lower symptoms on the CISS [46]. Divergent responses on symptom reporting on concussion-specific versus vision-specific surveys prevent an obvious explanation for the findings in the current student. Furthermore, previous studies on standard vision exams [28–30] and vestibular testing [32] reported differences in abnormal findings and increased prevalence rates across racial and SES groups, but those results do not align with the results from the current study. One exception was the VMS item, in which FRL group had a statistically significant greater symptom provocation score ($0.16 \pm .9$ vs. $0.06 \pm .6$); although the findings were no longer significant after accounting for multiple comparisons. These results did not appear to amount to any appreciable clinical significance as the differences at baseline were not only below clinical threshold, but within the SD range of the groups, and the effect size (Cohen's $d = .11$) was below the “small” threshold.

The debate regarding baseline testing for intra-individual changes versus normative comparison has long been discussed. While the most recent consensus statement did not suggest mandatory baseline testing, there are data to suggest that, for some clinical tools, intra-individual comparisons provide superior utility versus normative comparisons [67–69] and because various types of athletes [70–72], or athletes from varying backgrounds perform differently on testing [25,73,74]; however, this is not always the case [75–78]. Our model predicted less than 4% of the variance in K-D

test performance. These results indicate that performance is not well-predicted by race, sex, FRL status, and vision correction and provides support for baseline testing due to the inter-individual variability in our cohort.

As discussed previously, Title I schools provide funding to support their educational system but financial support does not always reach the sports medicine/athletic training departments on campus. So, while schools may appreciate the added benefit of baseline testing, it may not always be feasible. Fortunately, tools like the VOMS are publicly available and do not require a fee for use. However, there are always costs associated with baseline administrations, and while some tests are free, they are time intensive for staff members and students, and IT infrastructure necessary to securely and easily access medical records should be considered.

4.1 Limitations

First, there is no perfect method to determine individuals' SES, especially in a high school population, and thus, our findings are based on whether the student-athlete attended a school that met requirements for Title I or non-Title I status. This proxy for SES is widely utilized in educational research, but unfortunately, it is possible that some individual participants may belong to a different socioeconomic class that typically would not coincide with the need for FRL or Title I classification (e.g., affluent student at FRL school). However, we believe the quantity of those individuals is likely minimal. Furthermore, our data only included White and Black participants. Individuals from other racial or ethnic backgrounds (e.g., Asian, Hispanic/Latino) should be considered in future studies to add further diversity to test psychometrics and concussion literature. Lastly, there was potential for inter-rater errors in test implementation and measurement. This study implemented the VOMS and K-D tests, which can be more affordable and easily implemented concussion screening assessments. Future research should examine if results are consistent with advanced laboratory-based vestibular and oculomotor tools (i.e. eye tracking). Lastly, longitudinal studies are needed to determine modifying factors at post-concussion intervals (e.g., 3 days, 1 week, 1 month), including race/ethnicity and SES.

4.2 Conclusion:

As the clinical diagnosis and treatment of concussion actively continues to incorporate multisensory measures that include vestibular and vision-based oculomotor tools, a respect for sociodemographic diversity is warranted. With a higher probability of undiagnosed and uncorrected vision impairment, vestibular dysfunction, and saccadic eye tracking deficits likely to be more apparent as a consequence of poverty or health inequities, it is important that healthcare providers, especially those that diagnose and treat concussions, understand that the VOMS and K-D tests at baseline may be subject to sociodemographic factors of SES and race. Further, there is a greater likelihood of learning disabilities going unrecognized and undiagnosed among disadvantaged youth, which may adversely affect performance on assessments such as the K-D [56]. Clinically, these results further emphasize the need for individual baseline assessments and growing need for generating more research on tools such as the VOMS and K-D in racially and ethnically diverse groups to add diversity to concussion literature. Pre-injury or baseline testing is commonly practiced by athletic institutions; however, in the absence of baseline data, clinicians rely on normative data for comparisons. Unfortunately, the utilization of normative data instead of pre-injury comparisons may increase false-positive [79] or false-negative rates [69] as also shown in the use of cognitive tests. If a patient is of a minority racial group or of low SES, normative data may not be an accurate or equitable reflection of expected performance post-injury or at the time of medical clearance. This is critical given the wide array of SES and racially diverse backgrounds that engage in adolescent sport activity. It is recommended that when making interpretations on the potential clinical diagnosis of concussion using these tools, SES and race should be considered if comparing to norm-referenced data.

Funding: This study was supported by a University Research Council grant from Youngstown State University.

Acknowledgements: We would like to acknowledge all of the high school athletic trainers, and Master of Athletic Training Students affiliated with Youngstown State University for their dedication and assistance with administering all baseline measures.

Conflicts of Interest: The authorship team has no conflicts of interest to report for this work.

References

- [1] Harmon KG, Clugston JR, Dec K, Hainline B, Herring S, Kane SF, et al. American Medical Society for Sports Medicine position statement on concussion in sport. *Br J Sports Med* 2019;53:213–25.
- [2] Giza C, Hovda D. The new neurometabolic cascade of concussion. *Neurosurgery* 2014;75 Suppl 4:S24-33.
- [3] McCrory P, Meeuwisse W, Dvořák J, Aubry M, Bailes J, Broglio S, et al. Consensus statement on concussion in sport-the 5th international conference on concussion in sport held in Berlin, October 2016. *Br J Sports Med* [Internet] 2017;51:838–47. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28446457>
- [4] Dobson JL, Yarbrough MB, Perez J, Evans K, Buckley T. Sport-Related Concussion Induces Transient Cardiovascular Autonomic Dysfunction. *Am J Physiol - Regul Integr Comp Physiol* 2017;312:575–84.
- [5] Master CL, Scheiman M, Gallaway M, Goodman A, Robinson RL, Master SR, et al. Vision Diagnoses Are Common After Concussion in Adolescents. *Clin Pediatr (Phila)* 2016;55:260–7.
- [6] Murray NG, Szekely B, Moran R, Ryan G, Powell D, Munkasy BA, et al. Concussion history associated with increased postural control deficits after subsequent injury. *Physiol Meas* 2019;40:024001.
- [7] Goldberger J, Wheeler G, Sydenstricker E. Pellagra Incidence in Relation to Sex, Age, Season, Occupation, and “Disabling Sickness” in Seven Cotton-Mill Villages of South Carolina during 1916. *Public Health Rep* 1920;35:1650–64.
- [8] Goldberger J, Wheeler G, Sydenstricker E. A Study of the Relation of Family Income and Other Economic Factors to Pellagra Incidence in Seven Cotton-Mill Villages of South Carolina in 1916. *Public Health Rep* 1920;35:2673–714.
- [9] Houck Z, Asken B, Clugston J, Perlstein W, Bauer R. Socioeconomic Status and Race Outperform Concussion History and Sport Participation in Predicting Collegiate Athlete Baseline Neurocognitive Scores. *J Int Neuropsychol Soc* 2018;24:1–10.
- [10] Post E, Winterstein AP, Hetzel SJ, Lutes B, McGuine TA. School and community socioeconomic status and access to athletic trainer services in Wisconsin secondary schools. *J Athl Train* 2019;54:177–81.
- [11] Kroshus E, Rivara FP, Whitlock KB, Herring SA, Chrisman SPD. Disparities in Athletic Trainer Staffing in Secondary School Sport. *Clin J Sport Med* 2017;27:542–7.
- [12] Copley M, Jimenez N, Kroshus E, Chrisman SPD. Disparities in Use of Subspecialty Concussion Care Based on Ethnicity. *J Racial Ethn Heal Disparities* 2020;
- [13] Schneier AJ, Shields BJ, Hostetler SG, Xiang H, Smith GA. Incidence of pediatric traumatic brain injury and associated hospital resource utilization in the United States. *Pediatrics* 2006;118:483–92.
- [14] Information A. FACT SHEET : Supplement-not-Supplant under Title I of the Every Student Succeeds Act. 2016 [cited 2020 Jun 1];1–7. Available from: <https://www.ed.gov/news/press-releases/fact-sheet-supplement-not-supplant-under-title-i-every-student-succeeds-act>
- [15] Number and percentage of public school students eligible for free or reduced-price lunch, by

- state: Selected years, 2000-01 through 2015-16 [Internet]. [cited 2020 Jun 1]. Available from: https://nces.ed.gov/programs/digest/d17/tables/dt17_204.10.asp
- [16] Currie J, Thomas D. Medical care for children: public insurance, private insurance, and racial differences in utilization. *J Hum Resour* 1995;30:135–62.
- [17] Lyons TW, Miller KA, Miller AF, Mannix R. Racial and ethnic differences in emergency department utilization and diagnosis for sports-related head injuries. *Front Neurol* 2019;10.
- [18] Houck ZM, Asken BM, Bauer RM, Kontos AP, Mccrea MA, Mcallister TW, et al. Multivariate Base Rates of Low Scores and Reliable Decline on ImPACT in Healthy Collegiate Athletes Using CARE Consortium Norms. *J Int Neuropsychol Soc* 2019;25:961–71.
- [19] Kontos AP, Elbin RJ, Covassin T, Larson E. Exploring Differences in Computerized Neurocognitive Concussion Testing Between African American and White Athletes. *Arch Clin Neuropsychol* 2010;25:734–44.
- [20] O'Connor KL, Dain Allred C, Cameron KL, Campbell DE, D'Lauro CJ, Houston MN, et al. Descriptive Analysis of a Baseline Concussion Battery Among U.S. Service Academy Members: Results from the Concussion Assessment, Research, and Education (CARE) Consortium. *Mil Med* 2018;183:e580–90.
- [21] Semenov YR, Bigelow RT, Xue Q-L, Lac S Du, Agrawal Y. Association Between Vestibular and Cognitive Function in U.S. Adults: Data From the National Health and Nutrition Examination Survey. *Journals Gerontol Ser A Biol Sci Med Sci* 2016;71:243–50.
- [22] Norheim N, Kissinger-Knox A, Cheatham M, Webbe F. Performance of college athletes on the 10-item word list of SCAT5. *BMJ Open Sport Exerc Med* 2018;4:e000412.
- [23] Wallace J, Covassin T, Moran R, Deitrick JM. Factors Contributing to Disparities in Baseline Neurocognitive Performance and Concussion Symptom Scores Between Black and White Collegiate Athletes. *J Racial Ethn Heal Disparities* 2018;5:894–900.
- [24] Holmes L, Tworig J, Casini J, Morgan I, O'Brien K, Oceanic P, et al. Implication of Socio-Demographics on Cognitive-Related Symptoms in Sports Concussion Among Children. *Sport Med - Open* 2016;2.
- [25] Rexrode BL, Armstrong JL, Hallberg CT, Copeland BW, Bradney DA, Bowman TG. The effects of socioeconomic status on baseline neurocognitive testing scores. *Appl Neuropsychol Child* 2019;0:1–6.
- [26] Galetta KM, Barrett J, Allen M, Madda F, Delicata D, Tennant a. T, et al. The King-Devick test as a determinant of head trauma and concussion in boxers and MMA fighters. *Neurology [Internet]* 2011;76:1456–62. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21288984>
- [27] Mucha A, Collins MW, Elbin RJ, Furman JM, Troutman-Enseki C, DeWolf RM, et al. A Brief Vestibular/Ocular Motor Screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med [Internet]* 2014;42:2479–86. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25106780>
- [28] Frick DJ, Olitsky SE, Campbell A, Glaros AG. Ocular problems in low-income and minority children. *J Pediatr Ophthalmol Strabismus* 2013;50:363–8.
- [29] Logan NS, Shah P, Rudnicka AR, Gilmartin B, Owen CG. Childhood ethnic differences in ametropia and ocular biometry: the Aston Eye Study. *Ophthalmic Physiol Opt* 2011;31:550–8.
- [30] Ying GS, Maguire MG, Cyert LA, Ciner E, Quinn GE, Kulp MT, et al. Prevalence of Vision

Disorders by Racial and Ethnic Group among Children Participating in Head Start. *Ophthalmology* 2014;121:630–6.

- [31] Vitale S, Cotch MF, Sperduto RD. Prevalence of visual impairment in the United States. *J Am Med Assoc* 2006;295:2158–63.
- [32] Agrawal Y, Carey JP, Della Santina CC, Schubert MC, Minor LB. Disorders of balance and vestibular function in US adults: Data from the National Health and Nutrition Examination Survey, 2001–2004. *Arch Intern Med* 2009;169:938–44.
- [33] Scheiman M, Gallaway M, Frantz K a., Peters RJ, Hatch S, Cuff M, et al. Nearpoint of Convergence: Test Procedure, Target Selection, and Normative Data. *Optom Vis Sci* 2003;80:214–25.
- [34] Moran RN, Covassin T, Elbin RJ, Gould D, Nogle S. Reliability and Normative Reference Values for the Vestibular/Ocular Motor Screening (VOMS) Tool in Youth Athletes. *Am J Sports Med* 2018;46:1475–80.
- [35] Moran RN, Covassin T. King-Devick test normative reference values and internal consistency in youth football and soccer athletes. *Scand J Med Sci Sports* 2018;1–5.
- [36] Nagelkerke NJD. A note on a general definition of the coefficient of determination. *Biometrika* [Internet] 1991 [cited 2020 Sep 14];78:691–2. Available from: <https://academic.oup.com/biomet/article/78/3/691/256225>
- [37] Heick JD, Bay C, Dompier TP, Valovich McLeod TC, Leong D. The Psychometric Properties of the King–Devick Test and the Influence of Age and Sex in Healthy Individuals Aged 14 to 24 Years. *Athl Train Sport Heal Care* [Internet] 2016;8:222–9. Available from: <http://www.healio.com/doiresolver?doi=10.3928/19425864-20160509-01>
- [38] King-Devick Test®. Sideline Concussion Screening | King-Devick Technologies, Inc. [Internet]. 2018 [cited 2018 Oct 11]. Available from: <https://kingdevicktest.com/products/concussion/sideline-concussion-screening/>
- [39] Ganz ML, Xuan Z, Hunter DG. Prevalence and Correlates of Children’s Diagnosed Eye and Vision Conditions. *Ophthalmology* 2006;113:2298–306.
- [40] Ganz M, Xuan Z, Hunter DG. Patterns of eye care use and expenditures among children with diagnosed eye conditions. *J AAPOS* 2007;11:480–7.
- [41] Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: a Practical and Powerful Approach to Multiple Testing [Internet]. Vol. 57, *Journal of the Royal Statistical Society. Series B (Methodological)*. 1995. p. 289–300. Available from: http://www.stat.purdue.edu/~doerge/BIOINFORM.D/FALL06/Benjamini and Y FDR.pdf%5Cnhttp://enr.case.edu/ray_soumya/mlrg/controlling_fdr_benjamini95.pdf
- [42] Team RC. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2020.
- [43] Sherry NS, Fazio-Sumrok V, Sufrinko A, Collins MW, Kontos AP. Multimodal Assessment of Sport-Related Concussion. *Clin J Sport Med* 2019;00:1.
- [44] Felleman DJ, Van Essen DC. Distributed hierarchical processing in the primate cerebral cortex. *Cereb Cortex* 1991;1:1–47.
- [45] Galetta KM, Morganroth J, Moehringer N, Mueller B, Hasanaj L, Webb N, et al. Adding vision to concussion testing: A prospective study of sideline testing in youth and collegiate athletes. *J*

Neuro-Ophthalmology 2015;35:235–41.

- [46] Barnhardt C, Cotter SA, Mitchell GL, Scheiman M, Kulp MT. Symptoms in children with convergence insufficiency: Before and after treatment. *Optom Vis Sci* 2012;89:1512–20.
- [47] Kau CH, Wang J, Davis M. A Cross-Sectional Study to Understand 3D Facial Differences in a Population of African Americans and Caucasians. *Eur J Dent* 2019;13:485–96.
- [48] Phillips J, Cheever K, McKeon J, Tierney R. Effect of administration method on near point of convergence scores in a healthy, active, young adult population. *Int J Athl Ther Train* 2018;23:156–61.
- [49] SHANKAR S, EVANS MA, BOBIER WR. Hyperopia and Emergent Literacy of Young Children: Pilot Study. *Optom Vis Sci* 2007;84:1031–8.
- [50] Powers M, Grisham D, Riles P. Saccadic tracking skills of poor readers in high school. *Optometry* 2008;79:228–34.
- [51] Basch CE. Vision and the Achievement Gap Among Urban Minority Youth. *J Sch Health* 2011;81:599–605.
- [52] School-aged Vision: 6 to 18 Years of Age [Internet]. [cited 2020 Jun 1]. Available from: <https://www.aoa.org/patients-and-public/good-vision-throughout-life/childrens-vision/school-aged-vision-6-to-18-years-of-age>
- [53] Galetta KM, Chapman KR, Essis MD, Alosco ML, Gillard D, Steinberg E, et al. Screening Utility of the King-Devick Test in Mild Cognitive Impairment and Alzheimer Disease Dementia. *Alzheimer Dis Assoc Disord* 2017;31:152–8.
- [54] Sirin SRSR. Socioeconomic Status and Academic Achievement: A Meta-Analytic Review of Research. *Rev Educ Res* 2005;75:417–53.
- [55] Rosen ML, Amso D, McLaughlin KA. The role of the visual association cortex in scaffolding prefrontal cortex development: A novel mechanism linking socioeconomic status and executive function. Vol. 39, *Developmental Cognitive Neuroscience*. 2019.
- [56] Shifrer D, Muller C, Callahan R. Disproportionality and Learning Disabilities: Parsing Apart Race, Socioeconomic Status, and Language. *J Learn Disabil* [Internet] 2011;44:246–57. Available from: <http://10.0.4.153/0022219410374236>
- [57] Weise KK, Swanson MW, Penix K, Hale MH, Y DF. King-Devick and Pre-season Visual Function in Adolescent Athletes. *Optom Vis Sci* 2017;94:89–95.
- [58] Zhou W, Liu Y, Su M, Yan M, Shu H. Alternating-color words influence Chinese sentence reading: Evidence from neural connectivity. *Brain Lang* 2019;197.
- [59] Pyykkö J, Forssman L, Maleta K, Ashorn P, Ashorn U, Leppänen JM. Early development of visual attention in infants in rural Malawi. *Dev Sci* 2019;22.
- [60] Alsalaheen B, Haines J, Yorke A, Diebold J. King-Devick Test reference values and associations with balance measures in high school American football players. *Scand J Med Sci Sport* 2016;26:235–9.
- [61] Worts PR, Schatz P, Burkhart SO. Test Performance and Test-Retest Reliability of the Vestibular/Ocular Motor Screening and King-Devick Test in Adolescent Athletes During a Competitive Sport Season. *Am J Sports Med* 2018;46:2004–10.
- [62] Moser RS, Olek L, Schatz P. Gender Differences in Symptom Reporting on Baseline Sport

- Concussion Testing Across the Youth Age Span. *Arch Clin Neuropsychol* 2019;34:50–9.
- [63] Guo J, Yang J, Yi H, Singichetti B, Stavrinou D, Peek-Asa C. Differences in Postinjury Psychological Symptoms Between Collegiate Athletes With Concussions and Orthopedic Injuries. *Clin J Sport Med* 2018;0:1.
- [64] Cottle JE, Hall EE, Patel K, Barnes KP, Ketcham CJ. Concussion Baseline Testing: Preexisting Factors, Symptoms, and Neurocognitive Performance. *J Athl Train* 2017;52:77–81.
- [65] Benedict PA, Baner N V, Harrold GK, Moehringer N, Hasanaj L, Serrano LP, et al. Gender and age predict outcomes of cognitive, balance and vision testing in a multidisciplinary concussion center. *J Neurol Sci* [Internet] 2015;353:111–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25953343>
- [66] Covassin T, Schatz P, Swanik CB. Sex Differences in Neuropsychological Function and Post-Concussion Symptoms of Concussed Collegiate Athletes. *Neurosurgery* 2007;61:345–51.
- [67] Schmidt JD, Register-Mihalik JK, Mihalik JP, Kerr ZY, Guskiewicz KM. Identifying impairments after concussion: Normative data versus individualized baselines. *Med Sci Sport Exerc* 2012;44:1621–8.
- [68] Louey AG, Cromer JA, Schembri AJ, Darby DG, Maruff P, Makdissi M, et al. Detecting cognitive impairment after concussion: Sensitivity of change from baseline and normative data methods using the CogSport/Axon cognitive test battery. *Arch Clin Neuropsychol* 2014;29:432–41.
- [69] Schatz P, Robertshaw S. Comparing post-concussive neurocognitive test data to normative data presents risks for under-classifying “above average” athletes. *Arch Clin Neuropsychol* 2014;29:625–32.
- [70] Sandel N, Worts P, Burkhart S, Henry L. Comparison of baseline ImPACT performance in amateur motocross riders to football and basketball athletes. *Brain Inj* 2018;32:493–7.
- [71] Zimmer A, Picora K, Schuster D, Webbe F. Sport and team differences on baseline measures of sport-related concussion. *J Athl Train* 2013;48:659–67.
- [72] Katz BP, Kudela M, Harezlak J, McCrea M, McAllister T, Broglio SP, et al. Baseline Performance of NCAA Athletes on a Concussion Assessment Battery: A Report from the CARE Consortium. *Sports Med* 2018;48:1971–85.
- [73] Chrisman SPD, Harmon KG, Schmidt JD, Kaminski TW, Buckley TA, Kontos AP, et al. Impact of Factors that Affect Reading Skill Level on King–Devick Baseline Performance Time. *Ann Biomed Eng* 2018;47:2122–7.
- [74] Kontos AP, Sufrinko A, Elbin RJ, Puskar A, Collins MW. Reliability and Associated Risk Factors for Performance on the Vestibular/Ocular Motor Screening (VOMS) Tool in Healthy Collegiate Athletes. *Am J Sports Med* 2016;44:1400–6.
- [75] Tsushima WT, Oshiro R, Zimbra D. Neuropsychological test performance of Hawai'i high school athletes: Hawai'i ImPACT normative data. *Hawaii Med J* 2008;67:93–5.
- [76] Tsushima WT, Siu AM. Neuropsychological Test Performance of Hawai'i High School Athletes: Updated Hawai'i Immediate Post-Concussion Assessment and Cognitive Testing Data. *Hawai'i J Med Public Heal* 2014;73:208–11.
- [77] Shuttleworth-Edwards AB, Whitefield-Alexander VJ, Radloff SE, Taylor AM, Lovell MR. Computerized neuropsychological profiles of South African versus US athletes: a basis for

commentary on cross-cultural norming issues in the sports concussion arena. *Phys Sportsmed* 2009;37:45–52.

- [78] Shuttleworth-Edwards AB, Smith I, Radloff SE. Neurocognitive vulnerability amongst university rugby players versus noncontact sport controls. *J Clin Exp Neuropsychol* 2008;30:870–84.
- [79] Roebuck-Spencer TM, Vincent AS, Schlegel RE, Gilliland K. Evidence for added value of baseline testing in computer-based cognitive assessment. *J Athl Train* 2013;48:499–505.

Epub ahead of print